

Antiretroviral therapy, immunoreactivity and level of virus loading at the patients with co-infection HIV/tuberculosis

Makarov Pavel Viktorovich, M.D., assistant of faculty of infectious illnesses and epidemiology
Tver State Medical University

Summary. Russian federation concerns to number of the countries with high worsening tuberculosis epidemiological situation on HIV-infection. At the patients with HIV-infection of tuberculosis mortality reaches 43-89 %. The tuberculosis morbidity at the patients with HIV-infection were at 21-37 of time is higher, than at the people without HIV-infection, and in separate researches speak about risk of tuberculosis development at early stages of HIV-infection in 113 times.

The purpose of research - definition of antiretroviral therapy influence on immunoreactivity condition and level of virus loading at the patients with co-infection HIV/tuberculosis.

Totally 200 persons with a tuberculosis and HIV infection with a various mode of tuberculosis treatment were surveyed.

Under of antiretroviral therapy (ARVT) influence the CD4 lymphocytes level increased on 34-37 % in comparison with the patients, which did not received ARVT.

The antiretroviral therapy resulted to rise of CD4 lymphocytes percentage almost on 40 %. Antiretroviral therapy resulted to increase absolute and percentage CD4-lymphocytes, that is to activation of immune responses, which outcome was the difference in a level of HIV virus load for patients receiving ARVT in comparison with persons which do not receiving ones.

Keywords: tuberculosis, HIV-infection, immunoreactivity

Tuberculosis is one of the most often secondary diseases at the patients with HIV-infection causing a heavy defeat of various localization and being the basic reason of lethal outcome at the patients with AIDS [1, 3]. The patients having simultaneously a tuberculosis and a HIV-infection, concern to the most difficult contingent, both among the patients with HIV-infection, and among the patients with tuberculosis [4].

Presence of a HIV-infection is highest risk of tuberculosis development. Mortality from tuberculosis for patients with HIV-infection is more, than at the constant population of Russia consisting on the account, in 28 times. The influence of HIV-infection on epidemic process at the tuberculosis will be increased. By 2020 among the first time registered tuberculosis patients, HIV-infection can have up to 30 % of the patients as a whole on Russia, and in the separate subjects of Russian Federation - till 40-50 %. The tuberculosis morbidity of HIV-infected, in comparison with average parameters on Russia, among the constant population in 2012 was in 35 times more [2].

The purpose of research - definition of antiretroviral therapy influence on immunoreactivity condition and level of

virus loading at the patients with co - infection HIV/tuberculosis.

Materials and methods

In total 200 men were inspected. Group I - 100 patients with co-infection of lung tuberculosis and HIV, not receiving antiretroviral therapy (ARVT); group II - 100 patients with co-infection of lung tuberculosis and HIV, receiving antiretroviral therapy.

Results and their arguing

The standard procedure of inspection HIV-infected patients is the definition of the immune status, in particular of CD4 lymphocytes level.

Under of ARVT influence (tabl. 1) the CD4 lymphocytes level increased on 34,5 % (patients of 3-rd group) and 37,5 % (patients of 2 group) in comparison with the patients, which did not received ARVT (1-я group). For the 2 group patients the CD4 contents has made $373,8 \pm 13,6$ and $356,3 \pm 11,5$ cells/mm³ for the 3-rd group patients. The different is uncertain ($P > 0,05$).

Table 1. Immune status parameters at the patients with tuberculosis/ HIV/ on groups in absolute units

CD4-lymphocytes parameters	Groups of the inspected persons		P1
	Tuberculosis/ HIV group 1 (n=100)	Tuberculosis/HIV group 2 (n=100)	
CD4- lymphocytes parameters cells/mm ³	233,3±13,9	373,8±13,6	<0,001

The note: P1 - reliability of distinctions of parameters at the patients with tuberculosis/HIV 1- ű and 2 groups.

Taking into account that the absolute metrics of the immune status can considerably vary during even of day, the definition of CD4 lymphocytes level in percentage was carried out. In this case metrics considerably differed with a high degree of reliability for the patients of 1-st group on the one hand and patients of 2-rd groups, from other (tabl. 2).

The antiretroviral therapy resulted to rise of CD4 lymphocytes percentage almost on 40 %. However contents CD4

cells did not depend on the mode of antituberculosis therapy. The difference in percentage CD4 lymphocytes in group 2 and 3 has appeared statistically uncertain ($P > 0,05$).

One of main laboratory markers of HIV progressing now is the definition of HIV virus load (concentration of HIV RNA in blood). We pars metrics for the patients with a co-infection a HIV/tuberculosis in dependence of the antituberculosis therapy mode from spent or not spent ARVT (tabl. 3).

Table 2. Immune status parameters at the patients with HIV/tuberculosis on groups in %

CD4-lymphocytes parameters	Groups of the inspected persons		P1
	Tuberculosis/ HIV group 1 (n =100)	Tuberculosis/ HIV group 2 (n =100)	
CD4- lymphocytes parameters in %	13,1±1,5	21,3±1,3	<0,001

The note: P1 - reliability of distinctions of parameters at the patients with HIV/tuberculosis 1- ñ and 2 groups.

Table 3. Parameters of virus loading at the patients with co-infection HIV/tuberculosis on groups in absolute units

Parameters of virus loading	Groups of the inspected persons		P1 1-2
	Tuberculosis/ HIV group 1 (n =100)	Tuberculosis/ HIV group 2 (n =100)	
Virus loading, copy/мл	12000000±72000	68000±1200	<0,001

The note: P1 - reliability of distinctions of parameters at the patients with HIV/tuberculosis 1- ñ and 2 groups.

The essential decrease of virus loading under influence of antiretroviral therapy at the patients with lung tuberculosis receiving treatment on 1-st standard mode was established.

Conclusions

Antiretroviral therapy resulted to increase absolute and percentage CD4-lymphosites, that is to activation of immune responses, which outcome was the difference in a level of HIV virus load for patients receiving ARVT in comparison with persons which do not receiving ones practically in 176 time.

Literature:

1. Dolgova E.A., Al'vares Figeroa M. V., Lobasheva G. P. Definition of medicinal stability to rifampicine at the patients with tuberculosis which is taking place at a stage of secondary diseases of HIV-infection // Tub. i bolezni legkih. - 2014. - № 4. - S. 129-130.
2. Nechaeva O.B. A situation on a tuberculosis and HIV-infection in Russia // Tuberkulyoz i bolezni lyogkih. – 2014. № 6. – S. 9-15.
3. Pantelev A. M. Tuberculosis with medicinal stability MBT at patients with HIV-infection // Aktual. probl. i perspektivy protivotuberkuleznoj sluzhby v RF: Materialy I Kongressa Nacional'noj asociacii ftiziatrov. - SPb. - 2012. - S. 281-283.
4. Frolova O.G. SHCHukina I.V., Frolov G G. The mortality analysis from a tuberculosis, connected with HIV-infection // Tuberkulyoz i bolezni legkih. 2014; (7): 32-36.

SUMMARY

Antiretroviral therapy, immunoreactivity and level of virus loading at the patients with co-infection tuberculosis/ HIV
Makarov P.V., Makarov V.R.

Russian Federation concerns to number of the countries with high worsening tuberculosis epidemiological situation on HIV-infection. At the patients with HIV-infection of tuberculosis mortality reaches 43-89 %. The tuberculosis morbidity at the patients with HIV-infection were at 21-37 of time is higher, than at the people without HIV-infection, and in separate researches speak about risk of tuberculosis development at early stages of HIV-infection in 113 times.

The purpose of research - definition of antiretroviral therapy influence on immunoreactivity condition and level of virus loading at the patients with co-infection HIV/tuberculosis.

Totally 200 persons with a tuberculosis and HIV infection with a various mode of tuberculosis treatment were surveyed.

Under of antiretroviral therapy (ARVT) influence the CD4 lymphocytes level increased on 34-37 % in comparison with the patients, which did not received ARVT.

The antiretroviral therapy resulted to rise of CD4 lymphocytes percentage almost on 40 %. Antiretroviral therapy resulted to increase absolute and percentage CD4-lymphocytes, that is to activation of immune responses, which outcome was the difference in a level of HIV virus load for patients receiving ARVT in comparison with persons which do not receiving ones.

Key words: tuberculosis, HIV-infection, immunoreactivity

THE ITEMS OF INFORMATION of the AUTHORS:

Makarov Pavel Viktorovich – M.D., assistant of faculty of infectious illnesses and epidemiology. Tver state medical university

The home address: Russia, 170023, Tver, Lenin Avenue, house 42, sq. 35.

e-mail: global-create@yandex.ru